

**Overview****Useful For**

Used to detect the presence of *Candida albicans* and *Candida glabrata* DNA in vaginal samples as an aid to the diagnosis of vulvovaginal candidiasis in symptomatic women. Also used in the diagnosis of *Trichomonas vaginalis* infections.

**Method Name**

Nucleic acid amplification (NAA)

**NY State Available**

Yes

**Specimen****Specimen Type**

Swab

**Specimen Required**

**Collection Container/Tube:** APTIMA Multitest, vaginal, or unisex swab

**Specimen Volume:** One swab

**Collection Instructions:** Collect vaginal fluid sample using the Gen-Probe Aptima swab by contacting the swab to the lower third of the vaginal wall and rotating the swab for 10 to 30 seconds to absorb fluid. Immediately place the swab into the transport tube and carefully break the swab shaft against the side of the tube. Tightly screw on the cap. Submit one vaginal swab in APTIMA Multitest, vaginal, or unisex swab. Ship refrigerate.

**Specimen Minimum Volume**

One swab

**Reject Due To**

Hemolysis	NA
Lipemia	NA
Icterus	NA

# Test Definition: FNSVG

NuSwab Vaginitis (VG)

Other	grossly contaminated specimens, leaking or broken tube; inappropriate specimen transport conditions, including specimens received frozen; specimens received after prolonged delay (usually >72 hours); specimen in expired transport or incorrect transport device; specimens with inappropriate source for test requested; specimen with fixative or additives; Aptima urine transport; Aptima swab transport >30 days from collection; Aptima swab specimen without a swab; cleaning swab (white-shaft swab) in Aptima swab transport; any non-Gen-Probe swab submitted in Aptima transport device; transport device with multiple swabs; bloody or grossly mucoid specimens; bacterial swabs; specimen in ProbeTec UPT transport; ProbeTec Q-swabs
-------	--

## Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Swab	Refrigerated (preferred)	30 days	
	Ambient	30 days	

## Clinical and Interpretive

### Clinical Information

This test is intended to be used as an aid to the diagnosis of bacterial vaginosis (BV) in women with a clinical presentation consistent with this disorder. The BV test utilizes semiquantitative PCR analysis of the three most predictive marker organisms (*Atopobium vaginae*, BVAB-2, and *Megasphaera-1*) to generate a total score that correlates directly with the presence or absence of BV. In this test system, samples with a score of 0 to 1 are considered negative for BV, samples with a score of 3 to 6 are positive for BV, and samples with a score of 2 are indeterminate for BV.

### Reference Values

*Candida albicans*, NAA: Negative

*Candida glabrata*, NAA: Negative

*Trich vag* by NAA: Negative

## Performance

### PDF Report

No

### Day(s) Performed

Monday through Sunday

# Test Definition: FNSVG

NuSwab Vaginitis (VG)

## Report Available

5 to 8 days

## Performing Laboratory Location

LabCorp Burlington

## Fees and Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their Regional Manager. For assistance, contact [Customer Service](#).

### Test Classification

This test was developed and its performance characteristics determined by LabCorp. It has not been cleared or approved by the Food and Drug Administration. The FDA has determined that such clearance or approval is not necessary.

### CPT Code Information

87801

87798 x 3

87661

### LOINC® Information

Test ID	Test Order Name	Order LOINC Value
FNSVG	NuSwab Vaginitis (VG)	Not Provided

Result ID	Test Result Name	Result LOINC Value
Z4735	Atopobium vaginae	69565-0
Z4736	BVAB 2	69566-8
Z4737	Megasphaera 1	69567-6
Z4738	Candida albicans, NAA	69562-7
Z4739	Candida glabrata, NAA	69563-5
Z4740	Trich vag by NAA	62461-9